

Canine Atopic Dermatitis Immunotherapeutic

For Use in Dogs Only

Cytopoint™ aids in the reduction of clinical signs associated with atopic dermatitis in dogs.

Cytopoint™ is a ready-to-use, sterile liquid containing a caninized monoclonal antibody (mAb) against interleukin-31 (IL-31). IL-31 has been shown to induce pruritus in dogs in laboratory studies.

Cytopoint™ remains in circulation for several weeks. It exerts a therapeutic effect by binding to and neutralizing soluble IL-31, thus inhibiting pruritus and reducing skin lesions. Like other naturally-occurring antibodies and antibody-antigen complexes, elimination is via normal protein degradation pathways.


SAFETY: A field safety study has demonstrated that Cytopoint™ is well tolerated in dogs after subcutaneous injection. Adverse events occurred at a similar frequency between treated and placebo groups in a study of 245 canine patients presented to veterinary hospitals and diagnosed with atopic dermatitis. In this study, dogs were administered Cytopoint™ (1.0-3.3 mg/kg body weight) or placebo by the subcutaneous route on Days 0 and 28. Signs of patient discomfort on administration and adverse events occurred at a similar frequency between treatment groups. In addition, there was no clinically important difference in clinical pathology changes between groups following either dose. A wide variety of concomitant medications were safely used, including parasiticides, antibiotics, antifungals, corticosteroids, vaccines, immunotherapy, antihistamines and other antipruritics, such as oclacitinib and cyclosporine.

Cytopoint™ has also been demonstrated to be well-tolerated in dogs in a laboratory safety study in which 7 consecutive monthly subcutaneous injections were administered at doses of 3.3 mg/kg or 10 mg/kg body weight (12 dogs per group).

EFFECTIVENESS: Cytopoint™ has a demonstrated onset of efficacy within 1 day and maintains efficacy for at least 1 month following a single dose in the field study.


In a study of 211 canine patients presented to veterinary hospitals and diagnosed with atopic dermatitis, a single dose of Cytopoint™ (up to 2.0 mg/kg) or placebo was administered subcutaneously. The study confirmed a significant (P<0.05) increase in the frequency of treatment success compared to placebo for at least one month at a dose of 2.0 mg/kg. Treatment success measures were based on the owner assessment of pruritus (Visual Analog Scale; VAS) scores and veterinarian assessment of skin lesion scores (Canine Atopic Dermatitis Extent and Severity Index; CADESI-03).

**Treatment Success:
Veterinarian Assessment of CADESI-03
50% Reduction vs Baseline**



*Significantly different compared to placebo (P<0.05)
Fifty dogs were treated with 2.0 mg/kg of Cytopoint™ and 52 dogs were treated with placebo on Day 0.

**Treatment Success:
Owner Assessment of Pruritus VAS
20 mm Reduction vs Baseline**



*Significantly different compared to placebo (P<0.05)
Fifty dogs were treated with 2.0 mg/kg of Cytopoint™ and 52 dogs were treated with placebo on Day 0.

DIRECTIONS:

Cytopoint™ is available in 1-mL vials in four concentrations (10, 20, 30 or 40 mg). Administer Cytopoint™ at a minimum dose of 0.9 mg/lb (2 mg/kg) body weight. For convenience, the dosing tables below may be used as a guideline. Repeat administration every 4-8 weeks as needed in individual patients.

Dogs weighing < 5.0 lb (<2.3 kg):

Aseptically withdraw 0.09 mL/lb (0.2 mL/kg) from a single, 10-mg (Sky) vial and administer subcutaneously. Discard any remaining product.

Dogs weighing 5.0-40 lb (2.3-18.1 kg):

Aseptically withdraw the full volume of the appropriate vial according to the dosage table below and administer subcutaneously.

Dog Body Weight in Pounds	Dog Body Weight in Kilograms	Presentation			
		10 mg (Sky)	20 mg (Plum)	30 mg (Blush)	40 mg (Navy)
5.0-10	2.3-4.5	1 vial			
10.1-20	4.6-9.1		1 vial		
20.1-30	9.2-13.6			1 vial	
30.1-40	13.7-18.1				1 vial

Dogs weighing > 40 lb (18.1 kg):

A single dose requires a combination of vials, as outlined in the table below. Prior to administration, collect the number of vials indicated under each presentation according to the dog's body weight.

Aseptically draw the full volume from each vial into one syringe and administer subcutaneously as a single injection.

Dog Body Weight in Pounds	Dog Body Weight in Kilograms	Presentation			
		10 mg (Sky)	20 mg (Plum)	30 mg (Blush)	40 mg (Navy)
40.1-50	18.2-22.7	1 vial +			1 vial
50.1-60	22.8-27.2		1 vial +		1 vial
60.1-70	27.3-31.7			1 vial +	1 vial
70.1-80	31.8-36.3				2 vials
80.1-90	36.4-40.8	1 vial +			2 vials
90.1-100	40.9-45.4		1 vial +		2 vials
100.1-110	45.5-49.9			1 vial +	2 vials
110.1-120	50.0-54.4				3 vials
120.1-130	54.5-59.0	1 vial +			3 vials
130.1-140	59.1-63.5		1 vial +		3 vials
140.1-150	63.6-68.0			1 vial +	3 vials
150.1-160	68.1-72.6				4 vials
160.1-170	72.7-77.1	1 vial +			4 vials
170.1-180	77.2-81.6		1 vial +		4 vials
180.1-190	81.7-86.2			1 vial +	4 vials
190.1-200	86.3-90.7				5 vials

PRECAUTIONS:

- The product does not have a preservative. Each vial is for single use only. Inactivate unused contents.
- This product is intended for subcutaneous use only.
- Store upright at 2°-8°C. Prolonged exposure to higher temperatures and/or direct sunlight may adversely affect potency. Do not freeze.
- Use entire contents when first opened.
- Sterilized syringes and needles should be used to administer this product. Do not sterilize with chemicals because traces of disinfectant may inactivate the product.
- This product has not been tested in pregnant, lactating or breeding animals.
- Do not mix with other products.
- In case of human exposure, contact a physician.

For use by, or under the supervision of, a veterinarian.

Technical inquiries should be directed to Zoetis Inc. Veterinary Services, (888) 963-8471 (USA), (800) 461-0917 (Canada).

This product has been shown to be efficacious in treating healthy dogs with canine atopic dermatitis.

VLN 190/PCN 9580.02

zoetis

Zoetis Inc.
Kalamazoo, MI 49007, USA