

## Parvovirus Vaccine

Killed Virus

*Erysipelothrix*

*Rhusiopathiae*

*Leptospira Bratislava-*

*Canicola-Grippotyphosa-*

*Hardjo-Icterohaemorrhagiae-*

*Pomona Bacterin*

For use in swine only

FarrowSure® **GOLD B**

**PRODUCT DESCRIPTION:** FarrowSure GOLD B is for vaccination of healthy breeding swine as an aid in preventing reproductive failure caused by porcine parvovirus (PPV), erysipelas caused by *Erysipelothrix rhusiopathiae*, and leptospirosis caused by *Leptospira bratislava*, *L. canicola*, *L. grippotyphosa*, *L. hardjo*, *L. icterohaemorrhagiae* and *L. pomona*. An 18-week duration of immunity following vaccination has been demonstrated against erysipelas. FarrowSure GOLD B is a liquid preparation of porcine parvovirus grown on an established porcine cell line, a serum-free, clarified *E. rhusiopathiae* culture, and whole cell cultures of the 6 *Leptospira* serovars identified above. The antigens have been chemically inactivated and adjuvanted with 2 adjuvants, including Amphigen®, to enhance the immune response.

**DISEASE DESCRIPTION:** PPV and *Leptospira* are common agents of reproductive loss in swine. While infection with either of these pathogens may produce subclinical disease, infection with PPV during pregnancy may result in fetal resorption, fetal mummification and stillbirths. Infection by *Leptospira* during the second half of pregnancy may cause abortions and stillbirths; late-term abortions are the most important economic effect of leptospirosis. Leptospirosis caused by any of the serovars represented here cannot be clinically differentiated. Abortions may also occur in sows infected with *E. rhusiopathiae* during pregnancy.

**SAFETY AND EFFICACY:** The safety of FarrowSure GOLD B was demonstrated under field conditions in approximately 1,340 sows and gilts receiving single or multiple doses. No serious systemic or allergic reactions were observed following vaccination.

Efficacy of the fractions of FarrowSure GOLD B was demonstrated in controlled challenge-of-immunity and immunogenicity tests. Additionally, duration-of-immunity studies demonstrated that FarrowSure GOLD B protected pigs against *E. rhusiopathiae* challenge 18 weeks after the second vaccination.

### **DIRECTIONS:**

*General Directions:* Shake well. Aseptically administer 2 mL intramuscularly.

*Primary Vaccination:* Healthy swine should receive 2 doses 3–5 weeks apart with the second dose administered 2–4 weeks prior to breeding. Healthy gilts, however, should receive the second dose as near as possible to 14 days prior to breeding.

*Revaccination:* Revaccination with a single dose is recommended prior to subsequent breedings. Boars should be revaccinated semiannually.

Good animal husbandry and herd health management practices should be employed.

### **PRECAUTIONS:**

Store at 2°–7°C. Prolonged exposure to higher temperatures may adversely affect potency. Do not freeze.

Use entire contents when first opened.

Sterilized syringes and needles should be used to administer this vaccine.

Do not vaccinate within 21 days before slaughter.

Contains gentamicin as preservative.

Transient injection site swelling and/or inappetence may occur following vaccination.

As with many vaccines, anaphylaxis may occur after use. Initial antidote of epinephrine is recommended and should be followed with appropriate supportive therapy.

This product has been shown to be efficacious in healthy animals. A protective immune response may not be elicited if animals are incubating an infectious disease, are malnourished or parasitized, are stressed due to shipment or environmental conditions, are otherwise immunocompromised, or the vaccine is not administered in accordance with label directions.

Technical inquiries should be directed to Zoetis Inc. Veterinary Services, (888) 963-8471 (USA), (800) 461-0917 (Canada).

For veterinary use only

U.S. Veterinary License No. 190

Zoetis Inc.

Kalamazoo, MI 49007, USA